

# Proceedings of the Iowa Academy of Science

---

Volume 43 | Annual Issue

Article 27

---

1936

## Pathological Polyploidy in Seedlings of Corn and Sorghum

J. E. Sass

*Iowa State College*

Copyright ©1936 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

---

### Recommended Citation

Sass, J. E. (1936) "Pathological Polyploidy in Seedlings of Corn and Sorghum," *Proceedings of the Iowa Academy of Science*, 43(1), 160-160.

Available at: <https://scholarworks.uni.edu/pias/vol43/iss1/27>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact [scholarworks@uni.edu](mailto:scholarworks@uni.edu).

PATHOLOGICAL POLYPLOIDY IN SEEDLINGS OF  
CORN AND SORGHUM

J. E. SASS

The ethyl mercury phosphate present in a commercial seed dust produces a characteristic dwarfing of seedlings of cereals. The histological symptoms consist of hypertrophy of meristematic organs, and eventually of the apical meristem. These malformed organs consist of enormously enlarged, parenchymatous, multinucleate cells. The several nuclei in a cell vary greatly in size and chromatin content. The small "micronuclei" have less than the normal diploid complement of 20 chromosomes. The "giant nuclei" are polyploid, many of them containing well over 100 chromosomes. The multinucleate and polyploid conditions are brought about by incomplete, multipolar mitosis.

DEPARTMENT OF BOTANY,  
IOWA STATE COLLEGE,  
AMES, IOWA.